arXiv:1603.04284v1 [math.NA] 14 Mar 2016

SYMMETRIC KRONECKER PRODUCTS AND SEMICLASSICAL WAVE PACKETS

GEORGE A. HAGEDORN AND CAROLINE LASSER

Abstract. We investigate the iterated Kronecker product of a square matrix with itself and prove an invariance property for symmetric subspaces. This motivates the definition of an iterated symmetric Kronecker product and the derivation of an explicit formula for its action on vectors. We apply our result for describing a linear change in the matrix parametrization of semiclassical wave packets.

1. Introduction

The Kronecker product of matrices is known to be ubiquitous [VL00], and our aim here is to investigate the n-fold Kronecker product of a complex square matrix $M \in \mathbb{C}^{d\times d}$ with itself,

$$M^{n\otimes} \;=\; \underbrace{M\otimes\cdots\otimes M}_{n\,\text{times}}, \qquad n\in\mathbb{N},$$

and to apply our findings to the parametrization of semiclassical wave packets.

1.1. Two-fold symmetric Kronecker products.

In semidefinite programming (See for example [AHO98] or [Kle02, Appendix E].), the two-fold Kronecker product has notably occurred in combination with subspaces of a particular symmetry property. One considers the space

$$X_2 \;=\; \left\{x\in\mathbb{C}^{d^2} : x=\operatorname{vec}(X),\; X \;=\; X^t\in\mathbb{C}^{d\times d}\right\},$$

that contains those vectors that can be obtained by the row-wise vectorization of a complex symmetric $d\times d$ matrix. The dimension of the space X_2 is

$$L_2 \;=\; \tfrac{1}{2}\,d\,(d+1).$$

One can prove that this space is invariant under Kronecker products, in the sense that for all matrices $M\in\mathbb{C}^{d\times d}$, one has

$$(M\otimes M)\,x\in X_2, \quad \text{whenever } \; x\in X_2.$$

Now one uses the standard basis of $\mathbb{C}^{d^2}$ for constructing an orthonormal basis of the subspace X_2 and defines a corresponding sparse $L_2\times d^2$ matrix P_2 that has the basis vectors as its rows. The symmetric Kronecker product of M with itself is then the $L_2\times L_2$ matrix

$$S_2(M) \;=\; P_2\;(M\otimes M)\;P_2^*.$$

Date: March 15, 2016.
2010 *Mathematics Subject Classification.* 15A69, 15B10, 81Q20.
Key words and phrases. Kronecker product, symmetry, semiclassical wave packet.

1.2. **n-fold symmetric Kronecker products.** How does one extend this construction to symmetrizing n-fold Kronecker products? It is instructive to revisit the second order space in two dimensions and to write a vector $x \in X_2$ as

$$x \;=\; (x_{(2,0)},\, x_{(1,1)},\, x_{(1,1)},\, x_{(0,2)})^t.$$

This labelling uses the multi-indices $k = (k_1,\, k_2) \in \mathbb{N}^2$ with $k_1 + k_2 = 2$ in the redundant enumeration

$$\nu_2 \;=\; ((2,0),\, (1,1),\, (1,1),\, (0,2))^t\,.$$

This description allows for a straightforward extension to higher order n and dimension d. One works with a redundant enumeration ν_n of the multi-indices $k = (k_1,\, \ldots,\, k_d) \in \mathbb{N}^d$ with $k_1 + \cdots + k_d = n$, and defines

$$X_n \;=\; \left\{x \in \mathbb{C}^{d^n} : x_j = x_{j'} \;\text{ if }\; \nu_n(j) = \nu_n(j')\right\}.$$

The dimension of X_n equals the number of mult-indices in $\mathbb{N}^d$ of order n, that is the binomial coefficient

$$L_n \;=\; \binom{n+d-1}{n}.$$

And again, we can prove invariance in the sense that for all $M \in \mathbb{C}^{d\times d}$

$$M^{n\otimes n}\, x \in X_n, \quad \text{whenever } \; x \in X_n.$$

See Proposition 2. Then, we use the standard basis of $\mathbb{C}^{d^n}$ to build an orthonormal basis of X_n and assemble the corresponding sparse $L_n \times d^n$ matrix P_n. All this motivates the definition of the n-fold symmetric Kronecker product as

$$S_n(M) \;=\; P_n\, M^{n\otimes}\, P_n^*.$$

The matrix $S_n(M)$ is of size $L_n \times L_n$ and inherits structural properties as invertibility or unitarity from the matrix M. See Lemma 4.

Our main result Theorem 1 provides an explicit formula for the action of the matrix $S_n(M)$ in terms of multinomial coeffients and powers of the entries of the original matrix M. Labelling the components of a vector $y \in \mathbb{C}^{L_n}$ by multi-indices of order n, we obtain for all $k \in \mathbb{N}^d$ with $|k| = n$ that

$$\begin{aligned}&(S_n(M)\, y)_k\\ &= \frac{1}{\sqrt{k!}} \sum_{|\alpha_1|=k_1} \cdots \sum_{|\alpha_d|=k_d} \binom{k_1}{\alpha_1} \cdots \binom{k_d}{\alpha_d}\, m_1^{\alpha_1} \cdots m_d^{\alpha_d}\, \sqrt{(\alpha_1 + \cdots + \alpha_d)!}\; y_{\alpha_1+\cdots+\alpha_d},\end{aligned}$$

where $m_1,\, \ldots,\, m_d \in \mathbb{C}^d$ denote the row vectors of M. The summations weighted with multinomial coefficients stem from the n-fold Kronecker product $M^{n\otimes}$, whereas the square roots of the factorials originate in the orthonormalization of the row vectors of the matrix P_n.

1.3. **Application to semiclassical wave packets.** Our interest in n-fold symmetric Kronecker products emerged when studying linear changes in the parametrization of semiclassical wave packets. Semiclassical wave packets have first been proposed in [Hag85] as a multivariate non-isotropic generalization of the Hermite functions. See also [Hag98]. A family of semiclassical wave packets

$$\left\{\varphi_k[A,\, B] : k \in \mathbb{N}^d\right\}$$

is parametrized by two invertible matrices $A,\ B \in \mathrm{GL}(d,\mathbb{C})$ and forms an orthonormal basis of the Hilbert space of square integrable functions.

More recently, semiclassical wave packets have been also used for the numerical discretization of semiclassical quantum dynamics by E. Faou, V. Gradinaru, and C. Lubich [FGL09, Lub08]. The computationally demanding step of this method is the assembly of the Galerkin matrix for the potential function $V : \mathbb{R}^d \to \mathbb{R}$ according to

$$\langle \varphi_k[A,\,B], V\varphi_l[A,\,B]\rangle \;=\; \int_{\mathbb{R}^d} \overline{\varphi_k[A,\,B](x)}\, V(x)\, \varphi_l[A,\,B](x)\, dx.$$

In an effort to develop a new variant [BGH] that performs the quadrature for semiclassical wave packets with different parametrizing matrices A', B', the search began for the matrices describing a linear change in the parametrization. Our main application is Corollary 1 in Section 5.4. It gives an explicit formula in terms of an n-fold symmetric Kronecker product.

1.4. **Organization of the paper.** In the next Section,we start with some combinatorics for explicitly relating the lexicographic enumeration of multi-indices of order n with the redundant enumeration ν_n. Then we introduce the symmetric subspaces X_n in Section 3 and construct an orthonormal basis together with the corresponding matrix P_n. In Section 4, we define the n-fold symmetric Kronecker product and prove our main result Theorem 1. An introduction to semiclassical wave packets and the description of linear changes in their parametrization by symmetric Kronecker products is given in Section 5.

1.5. **Notation.** On some occasions we shall use the binomial coefficient

$$\binom{n}{j} \;=\; \frac{n!}{(n-j)!\,j!}, \qquad \text{for non-negative integers } n \geq j.$$

We write a multi-index $k \in \mathbb{N}^d$ as a row vector $k = (k_1,\,\ldots,\,k_d)$. We use the modulus $|k| = k_1 + \cdots + k_d$, and the multinomial coefficient

$$\binom{|k|}{k} \;=\; \frac{|k|!}{k_1!\,\cdots\,k_d!}, \qquad \text{for } \; k \in \mathbb{N}^d.$$

We adopt the convention that any multinomial coefficient with any negative argument is defined to be 0. We also use the kth power of a vector,

$$x^k \;=\; x_1^{k_1}\,\cdots\,x_d^{k_d}, \qquad x \in \mathbb{C}^d.$$

2. Combinatorics

2.1. **Reverse Lexicographic ordering.** First we enumerate the set multi-indices of order n in d dimensions,

$$\left\{k \in \mathbb{N}^d : |k| = n\right\}, \qquad n \in \mathbb{N},$$

in reverse lexicographic ordering and collect them as components of a formal vector denoted by ℓ_n. The length of the vector ℓ_n is the binomial coefficient

$$L_n \;=\; \binom{n+d-1}{n}$$

One can think of this in the following way [JHT]: The multi-indices k of order n in d dimensions are in a one-to-one correspondence with the sequences of n identical balls and $d-1$ identical sticks. The sticks partition the line into d bins into which

one can insert the n balls. (The first bin is to the left of all the sticks, and contains k_1 balls; the last bin is to the right of all the sticks, and contains k_d balls; for $2 \leq j \leq d-1$, the j^{th} bin is between sticks $j-1$ and j, and it contains k_j balls.) *E.g.*, the multi-index (3, 2, 0, 1) in four dimensions corresponds to

If all these objects were distinguishable, there would be $(n+d-1)!$ permutations, but since the balls are all identical, one must divide by $n!$, and since the sticks are all identical, one must divide by $(d-1)!$.

2.2. **A redundant enumeration.** Next we redundantly enumerate and collect multi-indices of modulus n in a vector ν_n of length d^n. We proceed recursively and set $\nu_0 = ((0, \dots, 0))^t$, $\nu_1 = (e_1^*, \dots, e_d^*)^t$, and

$$\nu_{n+1} \;=\; \operatorname{vec} \begin{pmatrix} \nu_n(1) + e_1^* & \dots & \nu_n(d^n) + e_1^* \\ \vdots & & \vdots \\ \nu_n(1) + e_d^* & \dots & \nu_n(d^n) + e_d^* \end{pmatrix}, \qquad n \geq 0,$$

where $e_1, \dots, e_d \in \mathbb{C}^d$ are the standard basis vectors of $\mathbb{C}^d$, and vec denotes the row-wise vectorization of a matrix.

For example, for $d = 2$, we have

$$\begin{aligned} \nu_2 &= ((2,0), (1,1), (1,1), (0,2))^t, \\ \nu_3 &= ((3,0), (2,1), (2,1), (1,2), (2,1), (1,2), (1,2), (0,3))^t. \end{aligned}$$

2.3. **A partition.** For relating the lexicographic and the redundant enumeration, we define the mapping

$$\sigma_n : \{1, \dots, L_n\} \to \mathcal{P}(\{1, \dots, d^n\})$$

so that for all $i \in \{1, \dots, L_n\}$ and $j \in \{1, \dots, d^n\}$ the following holds:

$$j \in \sigma_n(i) \iff \nu_n(j) = \ell_n(i).$$

For example, for $d = 2$, we have

$$\sigma_2(1) = \{1\}, \quad \sigma_2(2) = \{2, 3\}, \quad \sigma_2(3) = \{4\}$$

and

$$\sigma_3(1) = \{1\}, \quad \sigma_3(2) = \{2, 3, 5\}, \quad \sigma_3(3) = \{4, 6, 7\}, \quad \sigma_3(4) = \{8\}.$$

We observe the following partition property.

Lemma 1. *We have*

$$\#\sigma_n(i) \;=\; \binom{n}{\ell_n(i)}, \qquad i = 1, \dots, L_n,$$

and $\bigcup_{i=1,\dots,L_n} \sigma_n(i) = \{1, \dots d^n\}$*, where the union is pairwise disjoint.*

Proof. We first prove that we have a partition property. For any $j \in \{1, \ldots, d^n\}$ there exists $i \in \{1, \ldots, L_n\}$ so that $\nu_n(j) = \ell_n(i)$. So, we clearly have

$$\bigcup_{i=1,\ldots,L_n} \sigma_n(i) = \{1, \ldots, d^n\}.$$

Moreover, since $j \in \sigma_n(i) \cap \sigma_n(i')$ is equivalent to $\ell_n(i) = \ell_n(i')$, that is, $i = i'$, the union is disjoint.

For proving the claimed cardinality, we argue by induction. For $n = 0$, we have

$$\ell_0 \;=\; ((0, \ldots, 0)) \;=\; \nu_0, \qquad \sigma_0(1) = \{1^0\}, \qquad \#\sigma_0(1) = 1.$$

For the inductive step, we observe that in the redundant enumeration ν_n, the multi-index $k = (k_1, \ldots, k_d)$ can be generated from d possible entries in ν_{n-1},

$$(k_1 - 1,\ k_2, \ldots, k_d),\ \ldots,\ (k_1, \ldots, k_{d-1}, k_d - 1)$$

by adding $e_1^*, \ldots, e_d^*$, respectively. Of course, such an entry only belongs to ν_{n-1} if all its components are non-negative. For each of these indices with all entries non-negative, there is a unique number $j \in \{1, 2, \ldots, L_{n-1}\}$, such that $l_{n-1}(j)$ is the given index. If one of these indices has a negative entry, we define $j = -1$ and $\sigma_{n-1}(-1)$ to be the empty set, *i.e.*,

$$\sigma_{n-1}(-1) \;=\; \{\}, \quad \text{whose cardinality is 0.}$$

We list the d numbers defined this way as $i_1, \ldots, i_d$, and note that all the positive values in this list must be distinct. Then,

$$\begin{aligned}
\#\sigma_n(i) &= \sum_{m=1}^{d} \#\sigma_{n-1}(i_m) \\
&= \sum_{m=1}^{d} \binom{n-1}{k_1, \ldots, k_m - 1, \ldots, k_d} \\
&= \sum_{m=1}^{d} \begin{cases} \frac{(n-1)!}{k_1! \cdots (k_m-1)! \cdots k_d!} & k_m > 0 \\ 0 & k_m = 0 \end{cases} \\
&= \frac{(n-1)!\,(k_1 + \cdots + k_d)}{k_1! \cdots k_d!} \\
&= \binom{n}{k}.
\end{aligned}$$

□

3. Symmetric subspaces

We next analyze the symmetric spaces

$$X_n \;=\; \left\{ x \in \mathbb{C}^{d^n} : x_j = x_{j'} \ \text{ if } \ \nu_n(j) = \nu_n(j') \right\}, \qquad n \in \mathbb{N}.$$

We have $X_0 = \mathbb{C}$ and $X_1 = \mathbb{C}^d$, whereas X_n is a proper subset of $\mathbb{C}^{d^n}$ for $n \geq 2$.

For example, for $d = 2$,

$$\begin{aligned} X_2 &= \left\{x \in \mathbb{C}^4 : x_2 = x_3\right\}, \\ X_3 &= \left\{x \in \mathbb{C}^8 : x_2 = x_3 = x_5,\ x_4 = x_6 = x_7\right\}. \end{aligned}$$

Any vector $x \in X_n$ has d^n components, but the components that correspond to the same multi-index in the redundant enumeration $\nu_n(1), \ldots, \nu_n(d^n)$ have the same value. Hence, at most L_n components of $x \in X_n$ are different. They may be labelled by the multi-indices $\ell_n(1), \ldots, \ell_n(L_n)$, and we often refer to them by

$$x_{\ell_n(i)}, \qquad i = 1, \ldots, L_n.$$

The symmetric subspace X_2 has also been considered in [VLV15], and we next relate the alternative second order constructions to ours.

3.1. **The second order space.** The second order subspace

$$X_2 = \left\{x \in \mathbb{C}^{d^2} : x_j = x_{j'} \text{ if } \nu_2(j) = \nu_2(j')\right\}$$

can also be described in terms of matrices. Since

$$\nu_2 = \operatorname{vec} \begin{pmatrix} e_1^* + e_1^* & \ldots & e_d^* + e_1^* \\ \vdots & & \vdots \\ e_1^* + e_d^* & \ldots & e_d^* + e_d^* \end{pmatrix},$$

we may write

$$X_2 = \left\{x \in \mathbb{C}^{d^2} : x = \operatorname{vec}(X),\ X = X^t \in \mathbb{C}^{d\times d}\right\}.$$

Alternatively, as in [VLV15, §2.3], one may permute the standard basis vectors $e_1, \ldots, e_{d^2} \in \mathbb{C}^{d^2}$ according to the $d^2 \times d^2$ permutation matrix

$$\Pi_{dd} = \left(e_{1+0\cdot d}, e_{1+1\cdot d}, \ldots, e_{1+(d-1)\cdot d}, \ldots, e_{d+0\cdot d}, e_{d+1\cdot d}, \ldots, e_{d^2}\right)$$

and characterize the symmetric subspace as

$$X_2 = \left\{x \in \mathbb{C}^{d^2} : \Pi_{dd}\, x = x\right\}.$$

This way of writing the second order subspace corresponds to the definition of symmetric tensor spaces by permutations as given in [Hack12, Chapter 3.5].

3.2. **Relation between the subspaces.** Due to the recursive definition of the redundant multi-index enumeration, the symmetric subspaces of neighboring order can be related to each other as follows.

Lemma 2. *The symmetric subspace X_{n+1} is contained in the d-ary Cartesian product of the symmetric subspace X_n,*

$$X_{n+1} \subseteq X_n \times \cdots \times X_n, \qquad n \in \mathbb{N}.$$

Proof. We decompose a vector

$$x = (x^{(1)}, \ldots, x^{(d)})^t \in X_{n+1}$$

into d subvectors with d^n components each. The d^{n+1} components of x can be labelled by the multi-indices

$$\nu_n(1) + e_1^*, \ldots, \nu_n(d^n) + e_1^*, \ldots, \nu_n(1) + e_d^*, \ldots, \nu_n(d^n) + e_d^*,$$

so that the components of the subvector $x^{(m)}$, $m = 1, \ldots, d$, can be labelled by

$$\nu_n(1) + e_m^*, \ \ldots, \ \nu_n(d^n) + e_m^*.$$

Hence, $x_j^{(m)} = x_{j'}^{(m)}$ if $\nu_n(j) = \nu_n(j')$, and $x^{(m)} \in X_n$ for all $m = 1, \ldots, d$. □

The two-dimensional example $X_1 = \mathbb{C}^2$ and $X_2 = \{x \in \mathbb{C}^4 : x_2 = x_3\}$ shows that the inclusion of Lemma 2 is in general not an equality.

3.3. **An orthonormal basis.** We now use the standard basis of $\mathbb{C}^{d^n}$ to construct an orthonormal basis of X_n.

Lemma 3. *Let $e_1, \ldots, e_{d^n}$ be the standard basis vectors of $\mathbb{C}^{d^n}$, and define the vectors*

$$p_i \ = \ \frac{1}{\sqrt{\#\sigma_n(i)}} \sum_{j \in \sigma_n(i)} e_j, \qquad i = 1, \ldots, L_n.$$

Then, $\{p_1, \ldots, p_{L_n}\}$ forms an orthonormal basis of the space X_n.

Proof. For all $i = 0, \ldots, L_n$ and $j, j' = 1, \ldots, d^n$, we have

$$(p_i)_j \ = \ \begin{cases} (\#\sigma_n(i))^{-1/2}, & \text{if } j \in \sigma_n(i), \\ 0, & \text{otherwise.} \end{cases}$$

Since $j \in \sigma_n(i)$ if and only if $\nu_n(j) = \ell_n(i)$, we have

$$(p_i)_j \ = \ (p_i)_{j'} \quad \text{if} \quad \nu_n(j) = \nu_n(j'),$$

and thus $p_i \in X_n$. We also observe, that for all $i, i' = 1, \ldots, L_n$,

$$\langle p_i, \, p_{i'} \rangle \ = \ \frac{1}{\sqrt{\#\sigma_n(i) \cdot \#\sigma_n(i')}} \sum_{j \in \sigma_n(i)} \sum_{j' \in \sigma_n(i')} \langle e_j, \, e_{j'} \rangle \ = \ \delta_{i,i'}.$$

Hence, the vectors $p_1, \ldots, p_{L_n}$ are orthonormal. Moreover, for all $x \in X_n$, we have

$$\langle p_i, \, x \rangle \ = \ \frac{1}{\sqrt{\#\sigma_n(i)}} \sum_{j \in \sigma_n(i)} \langle e_j, \ x \rangle \ = \ \sqrt{\#\sigma_n(i)}\, x_{\ell_n(i)},$$

and therefore

$$\begin{aligned} x \ &= \ \sum_{j=1}^{d^n} \langle e_j, \, x \rangle \, e_j \ = \ \sum_{i=1}^{L_n} \sum_{j \in \sigma_n(i)} \langle e_j, \, x \rangle \, e_j \ = \ \sum_{i=1}^{L_n} x_{\ell_n(i)} \sqrt{\#\sigma_n(i)}\, p_i \\ &= \ \sum_{i=1}^{L_n} \langle p_i, \, x \rangle \, p_i. \end{aligned}$$

□

3.4. **An orthonormal matrix.** The orthonormal basis vectors $p_1, \ldots, p_{L_n} \in X_n$ allow us to define the sparse rectangular $L_n \times d^n$ matrix

$$P_n \ = \ \begin{pmatrix} p_1^* \\ \vdots \\ p_{L_n}^* \end{pmatrix}$$

that has the L_n basis vectors as its rows. For example, for $d = 2$, we have

$$P_2 = \begin{pmatrix} e_1^* \\ \frac{1}{\sqrt{2}}(e_2^* + e_3^*) \\ e_4^* \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

$$P_3 = \begin{pmatrix} e_1^* \\ \frac{1}{\sqrt{2}}(e_2^* + e_3^* + e_5^*) \\ \frac{1}{\sqrt{2}}(e_4^* + e_6^* + e_7^*) \\ e_8^* \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

We summarize some properties of the matrix P_n and of its adjoint P_n^* and calculate explicit formulas for their actions on vectors.

Proposition 1. *The $L_n \times d^n$ matrix P_n and its adjoint P_n^* satisfy*

$$P_n P_n^* = \mathrm{Id}_{L_n \times L_n}, \qquad \mathrm{range}(P_n^*) = X_n.$$

Moreover, for all $x \in X_n$,

$$(P_n x)_i = \sqrt{\binom{n}{\ell_n(i)}}\, x_{\ell_n(i)}, \qquad i = 1, \ldots, L_n,$$

and for all $y \in \mathbb{C}^{L_n}$,

$$(P_n^* y)_{\ell_n(i)} = 1 \Big/ \sqrt{\binom{n}{\ell_n(i)}}\; y_i, \qquad i = 1, \ldots, L_n.$$

In particular,

$$P_n^* P_n x = x, \quad \text{whenever } x \in X_n.$$

Proof. The two properties $P_n P_n^* = \mathrm{Id}_{L_n \times L_n}$ and $\mathrm{range}(P_n^*) = X_n$ equivalently say, that the row vectors of P_n build an orthonormal basis of X_n.

For any $y \in \mathbb{C}^{L_n}$, the vector $P_n^* y$ is a linear combination of the vectors $p_1, \ldots, p_{L_n}$ and therefore in X_n. Labelling its components by $\ell_n(1), \ldots, \ell_n(L_n)$, we obtain

$$(P_n^* y)_{\ell_n(i)} = \sum_{i'=1}^{L_n} (p_{i'})_{\ell_n(i)}\, y_{i'} = \frac{1}{\sqrt{\#\sigma_n(i)}}\, y_i, \qquad i = 1, \ldots, L_n.$$

For $x \in X_n$ and $i = 1, \ldots, L_n$, we obtain

$$\begin{aligned} (P_n x)_i &= \sum_{j=1}^{d^n} (p_i)_{\nu_n(j)}\, x_{\nu_n(j)} = \frac{\#\sigma_n(i)}{\sqrt{\#\sigma_n(i)}}\, x_{\ell_n(i)} \\ &= \sqrt{\#\sigma_n(i)}\, x_{\ell_n(i)}, \end{aligned}$$

since $\#\sigma_n(i)$ components of p_i do not vanish. In particular,

$$(P_n^* P_n x)_{\ell_n(i)} = \frac{1}{\sqrt{\#\sigma_n(i)}}\, (P_n x)_i = x_{\ell_n(i)}.$$

□

4. Symmetric Kronecker products

4.1. **Iterated Kronecker products.** We next investigate the action of an n-fold Kronecker product on the symmetric subspace X_n, $n \in \mathbb{N}$.

Proposition 2. *Let $M \in \mathbb{C}^{d\times d}$, and denote by $m_1, \dots, m_d \in \mathbb{C}^d$ the row vectors of the matrix M. Then,*

$$M^{n\otimes}x \in X_n, \qquad \text{whenever } x \in X_n.$$

Moreover, for all $k \in \mathbb{N}^d$ with $|k| = n$, and all $x \in X_n$,

$$\big(M^{n\otimes}x\big)_k \;=\; \sum_{|\alpha_1|=k_1} \cdots \sum_{|\alpha_d|=k_d} \binom{k_1}{\alpha_1} \cdots \binom{k_d}{\alpha_d} m_1^{\alpha_1} \cdots m_d^{\alpha_d}\; x_{\alpha_1+\cdots+\alpha_d},$$

where $\alpha_1, \dots, \alpha_d \in \mathbb{N}^d$.

Proof. For $n = 1$, we have $M^{n\otimes} = M$ and $X_n = \mathbb{C}^d$, and our formula reduces to usual matrix-vector multiplication written as

$$(Mx)_{e_k} \;=\; \sum_{j=1}^{d} m_j^{e_k}\, x_{e_j}, \qquad k = 1, \dots, d.$$

For the inductive step, we consider $x = (x^{(1)}, \dots, x^{(d)}) \in X_{n+1}$ decomposed as in Lemma 2 with $x^{(j)} \in X_n$. We compute

$$\begin{aligned} M^{(n+1)\otimes}x \;&=\; \begin{pmatrix} m_{11}\, M^{n\otimes} & \dots & m_{1d}\, M^{n\otimes} \\ \vdots & & \vdots \\ m_{d1}\, M^{n\otimes} & \dots & m_{dd}\, M^{n\otimes} \end{pmatrix} \begin{pmatrix} x^{(1)} \\ \vdots \\ x^{(d)} \end{pmatrix} \\ &=\; \begin{pmatrix} m_{11}\, M^{n\otimes}x^{(1)} + \cdots + m_{1d}\, M^{n\otimes}x^{(d)} \\ \vdots \\ m_{d1}\, M^{n\otimes}x^{(1)} + \cdots + m_{dd}\, M^{n\otimes}x^{(d)} \end{pmatrix}. \end{aligned}$$

By the inductive hypothesis, we have for all $j = 1, \dots, d$, that

$$m_{j1}\, M^{n\otimes}x^{(1)} + \cdots + m_{jd}\, M^{n\otimes}x^{(d)} \in X_n.$$

The components of these vectors can be labelled by $k \in \mathbb{N}^d$ with $|k| = n$, and we have

$$\begin{aligned} &\Big(m_{j1}M^{n\otimes}x^{(1)} + \cdots + m_{jd}\, M^{n\otimes}x^{(d)}\Big)_k \\ &= \sum_{|\alpha_1|=k_1} \cdots \sum_{|\alpha_d|=k_d} \binom{k_1}{\alpha_1} \cdots \binom{k_d}{\alpha_d} \Big(m_{j1} m_1^{\alpha_1} \cdots m_d^{\alpha_d}\, x^{(1)}_{\alpha_1+\cdots+\alpha_d} + \cdots \\ &\qquad\qquad + m_{jd}\, m_1^{\alpha_1} \cdots m_d^{\alpha_d}\, x^{(d)}_{\alpha_1+\cdots+\alpha_d}\Big). \end{aligned}$$

We now observe that

$$\begin{aligned}
&\sum_{|\alpha_j|=k_j} \binom{k_j}{\alpha_j} \left(m_j^{\alpha_j+e_1} x^{(1)}_{\alpha_1+\cdots+\alpha_d} + \cdots + m_j^{\alpha_j+e_d} x^{(d)}_{\alpha_1+\cdots+\alpha_d} \right) \\
&= \sum_{|\alpha_j-e_1|=k_j} \binom{k_j}{\alpha_j - e_1} m_j^{\alpha_j} x^{(1)}_{\alpha_1+\cdots+(\alpha_j-e_1)+\cdots+\alpha_d} + \cdots + \\
&\qquad\qquad \sum_{|\alpha_j-e_d|=k_j} \binom{k_j}{\alpha_j - e_d} m_j^{\alpha_j} x^{(d)}_{\alpha_1+\cdots+(\alpha_j-e_d)+\cdots+\alpha_d} \\
&= \sum_{|\alpha_j|=k_j+1} \binom{k_j+1}{\alpha_j} m_j^{\alpha_j} x_{\alpha_1+\cdots+\alpha_d}.
\end{aligned}$$

Therefore, for all $j = 1, \ldots, d$,

$$\begin{aligned}
&\left(m_{j1} M^{n\otimes} x^{(1)} + \cdots + m_{jd} M^{n\otimes} x^{(d)} \right)_k = \\
&\sum_{|\alpha_1|=k_1} \cdots \sum_{|\alpha_j|=k_j+1} \cdots \sum_{|\alpha_d|=k_d} \binom{k_1}{\alpha_1} \cdots \binom{k_j+1}{\alpha_j} \cdots \binom{k_d}{\alpha_d} m_1^{\alpha_1} \cdots m_d^{\alpha_d} x_{\alpha_1+\cdots+\alpha_d},
\end{aligned}$$

Altogether, we have proven for all $k \in \mathbb{N}^d$ with $|k| = n+1$, and all $x \in X_{n+1}$

$$\left(M^{(n+1)\otimes} x \right)_k = \sum_{|\alpha|=k_1} \cdots \sum_{|\alpha_d|=k_d} \binom{k_1}{\alpha_1} \cdots \binom{k_d}{\alpha_d} m_1^{\alpha_1} \cdots m_d^{\alpha_d} x_{\alpha_1+\cdots+\alpha_d}.$$

□

4.2. **Symmetric Kronecker products.** Having proven that n-fold Kronecker products leave the nth symmetric subspace invariant, we define the n-fold symmetric Kronecker product as follows:

Definition 1. *For $M \in \mathbb{C}^{d\times d}$ and $n \in \mathbb{N}$, we define the $L_n \times L_n$ matrix*

$$S_n(M) = P_n M^{n\otimes} P_n^*$$

and call it the n-fold symmetric Kronecker product *of the matrix M.*

The n-fold symmetric Kronecker product has useful structural properties.

Lemma 4. *The n-fold symmetric Kronecker product $S_n(M)$ of a matrix $M \in \mathbb{C}^{d\times d}$ satisfies $S_n(M)^* = S_n(M^*)$. If $M \in \mathrm{GL}(d,\mathbb{C})$, then*

$$S_n(M) \in \mathrm{GL}(L_n,\mathbb{C}) \quad \text{with} \quad S_n(M)^{-1} = S_n(M^{-1}).$$

In particular, if $M \in U(d)$, then $S_n(M) \in U(L_d)$.

Proof. Since $(M\otimes M)^* = M^*\otimes M^*$ and $(M^{n\otimes})^* = (M^*)^{n\otimes}$, we have

$$S_n(M^*) = P_n (M^{n\otimes})^* P_n^* = S_n(M^*).$$

If M is invertible, then $M\otimes M$ is invertible with $(M\otimes M)^{-1} = M^{-1}\otimes M^{-1}$. By Proposition 2,

$$(M^{n\otimes})^{-1} p_j \in X_n, \qquad j = 1, \ldots, L_n.$$

Proposition 1 yields $P_n^* P_n x = x$ for all $x \in X_n$ and

$$P_n p_j = e_j, \qquad j = 1, \ldots, L_n,$$

where $e_1, \ldots, e_{L_n} \in \mathbb{C}^{L_n}$ are the standard basis vectors. Therefore,

$$S_n(M)\, S_n(M^{-1})\, e_j \;=\; P_n\, M^{n\otimes}\, P_n^*\, P_n\, (M^{n\otimes})^{-1} p_j \;=\; P_n\, p_j \;=\; e_j,$$

that is,

$$S_n(M)\, S_n(M^{-1}) \;=\; \mathrm{Id}_{L_n \times L_n}.$$

□

4.3. **The main result.** The explicit formula of Proposition 2 for the n-fold Kronecker product also allows a detailed description of the n-fold symmetric Kronecker product.

Theorem 1. *Let $M \in \mathbb{C}^{d\times d}$, and denote by $m_1, \ldots, m_d \in \mathbb{C}^d$ the row vectors of the matrix M. Then, the n-fold symmetric Kronecker product satisfies*

$$\begin{aligned}&(S_n(M)\, y)_k \\ &= \frac{1}{\sqrt{k!}} \sum_{|\alpha_1|=k_1} \cdots \sum_{|\alpha_d|=k_d} \binom{k_1}{\alpha_1} \cdots \binom{k_d}{\alpha_d}\, m_1^{\alpha_1} \cdots m_d^{\alpha_d}\, \sqrt{(\alpha_1+\cdots+\alpha_d)!}\; y_{\alpha_1+\cdots+\alpha_d}.\end{aligned}$$

for all $y \in \mathbb{C}^{L_n}$ and $k \in \mathbb{N}^d$ with $|k| = n$.

Proof. By Proposition 1, we have $P_n^*\, y \in X_n$ and

$$(P_n^*\, y)_{\ell_n(i)} \;=\; 1 \Bigg/ \sqrt{\binom{n}{\ell_n(i)}}\; y_i, \qquad \text{for } i = 1, \ldots, L_n.$$

By Proposition 2, the n-fold Kronecker product leaves X_n invariant, and we have for all $k \in \mathbb{N}^d$ with $|k| = n$,

$$\begin{aligned}\big(M^{n\otimes}\, P_n^*\, y\big)_k \;&=\; \sum_{|\alpha_1|=k_1} \cdots \sum_{|\alpha_d|=k_d} \binom{k_1}{\alpha_1} \cdots \binom{k_d}{\alpha_d}\, m_1^{\alpha_1} \cdots m_d^{\alpha_d}\, (P_n^*\, y)_{\alpha_1+\cdots+\alpha_d} \\ &= \sum_{|\alpha_1|=k_1} \cdots \sum_{|\alpha_d|=k_d} \binom{k_1}{\alpha_1} \cdots \binom{k_d}{\alpha_d}\, m_1^{\alpha_1} \cdots m_d^{\alpha_d}\, \sqrt{\frac{(\alpha_1+\cdots+\alpha_d)!}{n!}}\; y_{\alpha_1+\cdots+\alpha_d}.\end{aligned}$$

By Proposition 1, we have

$$(P_n\, x)_i \;=\; \sqrt{\binom{n}{\ell_n(i)}}\; x_{\ell_n(i)}, \qquad \text{for } i = 1, \ldots, L_n,$$

so that

$$\begin{aligned}&\big(P_n\, M^{n\otimes}\, P_n^*\, y\big)_k \\ &= \sqrt{\frac{n!}{k!}} \sum_{|\alpha_1|=k_1} \cdots \sum_{|\alpha_d|=k_d} \binom{k_1}{\alpha_1} \cdots \binom{k_d}{\alpha_d} m_1^{\alpha_1} \cdots m_d^{\alpha_d} \sqrt{\frac{(\alpha_1+\cdots+\alpha_d)!}{n!}}\, y_{\alpha_1+\cdots+\alpha_d} \\ &= \frac{1}{\sqrt{k!}} \sum_{|\alpha_1|=k_1} \cdots \sum_{|\alpha_d|=k_d} \binom{k_1}{\alpha_1} \cdots \binom{k_d}{\alpha_d} m_1^{\alpha_1} \cdots m_d^{\alpha_d} \sqrt{(\alpha_1+\cdots+\alpha_d)!}\; y_{\alpha_1+\cdots+\alpha_d}.\end{aligned}$$

□

5. Application to semiclassical wavepackets

5.1. **Parametrizing Gaussians.** We consider two complex invertible matrices $A,\, B \in \mathrm{GL}(d,\, \mathbb{C})$ that satisfy the conditions

$$A^t B - B^t A \;=\; 0, \tag{1}$$

$$A^* B + B^* A \;=\; 2\,\mathrm{Id}_{d\times d}. \tag{2}$$

These two conditions imply that $B\,A^{-1}$ is a complex symmetric matrix such that its real part satisfies

$$\mathrm{Re}(B\,A^{-1}) \;=\; (A\,A^*)^{-1}$$

is a Hermitian and positive definite matrix, see [Hag80]. Let $\hbar > 0$ and define the multivariate complex-valued Gaussian function

$$\varphi_0[A,\, B] : \mathbb{R}^d \to \mathbb{C},$$

$$\varphi_0[A,\, B](x) \;=\; (\pi\,\hbar)^{-d/4}\,\det(A)^{-1/2}\,\exp\left(-\frac{\langle x,\, B\,A^{-1}\,x\rangle}{2\,\hbar}\right).$$

Then, $\varphi_0[A,\, B]$ is a square-integrable function, and the constant factor $\det(A)^{-1/2}$ ensures normalization according to

$$\int_{\mathbb{R}^d} |\varphi_0[A,\, B](x)|^2\; dx \;=\; 1.$$

Changing the parametrization by a unitary matrix changes the Gaussian function only by constant multiplicative factor of modulus one:

Lemma 5. *Let $A,\, B \in \mathrm{GL}(d,\, \mathbb{C})$ satisfy the conditions (1–2). Then, for all unitary matrices $U \in U(d)$, the matrices $A' = A\,U$ and $B' = B\,U$ also satisfy the conditions (1–2). Moreover,*

$$\varphi_0[A',\, B'] = \det(U)^{-1/2}\,\varphi_0[A,\, B].$$

Proof. We observe that

$$(A')^t B' - (B')^t A' \;=\; U^t (A^t B - BA) U \;=\; 0$$

$$(A')^* B' + (B')^* A' \;=\; U^* (A^* B + BA) U \;=\; 2\,\mathrm{Id}_{d\times d}.$$

and $B'\,(A')^{-1} = B\,U\,U^* A^{-1} = BA^{-1}$. Therefore,

$$\begin{aligned}\varphi_0[A',\, B'] \;&=\; (\pi\,\hbar)^{-d/4}\,\det(A\,U)^{-1/2}\,\exp\left(-\frac{\langle x,\, B\,A^{-1}\,x\rangle}{2\,\hbar}\right)\\ &=\; \det(U)^{-1/2}\,\varphi_0[A,\, B].\end{aligned}$$

□

5.2. **Semiclassical wave packets.** Following the construction of [Hag98], we consider $A,\, B \in \mathrm{GL}(d,\, \mathbb{C})$ satisfying the conditions (1–2) and introduce the vector of raising operators

$$\mathcal{R}[A,\, B] \;=\; \frac{1}{\sqrt{2\hbar}}\,(B^*\,x - iA^*(-i\hbar\nabla_x))$$

that consists of d components,

$$\mathcal{R}[A,\, B] \;=\; \begin{pmatrix}\mathcal{R}_1[A,\, B]\\ \vdots\\ \mathcal{R}_d[A,\, B]\end{pmatrix}.$$

The raising operator acts on Schwartz functions $\psi : \mathbb{R}^d \to \mathbb{C}$ as

$$(\mathcal{R}[A,\, B]\psi)\,(x) \;=\; \frac{1}{\sqrt{2\hbar}}\,(B^*\,x\,\psi(x) - iA^*(-i\hbar\nabla_x\psi)(x))\,, \qquad x \in \mathbb{R}^d.$$

Powers of the raising operator generate the semiclassical wave packets according to

$$\varphi_k[A,\, B] \;=\; \frac{1}{\sqrt{k!}}\mathcal{R}[A,B]^k\varphi_0[A,\, B], \qquad k \in \mathbb{N}^d,$$

where the kth power of the raising operator

$$\mathcal{R}[A,\, B]^k = \mathcal{R}_1[A,\, B]^{k_1}\,\cdots\,\mathcal{R}_d[A,B]^{k_d}$$

does not depend on the ordering, since the components commute with one another due to the compatibility conditions (1–2). The set

$$\left\{\varphi_k[A,\, B] : k \in \mathbb{N}^d\right\}$$

forms an orthonormal basis of the space of square-integrable functions $L^2(\mathbb{R}^d,\, \mathbb{C})$.

5.3. **Hermite polynomials.** By its construction, the kth semiclassical wave packet is a multivariate polynomial of degree $|k|$ times the Gaussian function, that is,

$$\varphi_k[A,\, B] \;=\; \frac{1}{\sqrt{2^{|k|}\,k!}}\,p_k[A]\,\varphi_0[A,\, B].$$

The polynomials $p_k[A]$ are determined by the matrix $A \in \mathrm{GL}(d,\, \mathbb{C})$ and satisfy the three-term recurrence relation

$$\left(p_{k+e_j}[A]\right)_{j=1}^{d} \;=\; \frac{2}{\sqrt{\hbar}}A^{-1}x\,p_k[A] - 2A^{-1}\overline{A}\left(k_j\,p_{k-e_j}[A]\right)_{j=1}^{d}.$$

Whenever the parameter matrix A has all entries real, $A \in \mathrm{GL}(d,\, \mathbb{R})$, then the polynomials factorize according to

$$p_k[A](x) \;=\; \prod_{j=1}^{d} H_{k_j}\left(\tfrac{1}{\sqrt{\hbar}}(A^{-1}x)_j\right), \qquad x \in \mathbb{R}^d,$$

where H_n is the nth Hermite polynomial, $n \in \mathbb{N}$, defined by the univariate three-term recurrence relation

$$H_{n+1}(y) \;=\; 2\,y\,H_n(y) \;-\; 2\,n\,H_{n-1}(y), \qquad y \in \mathbb{R}.$$

The real parameter case, however, is rather expectional when using semiclassical wave packets for their key application in molecular quantum dynamics. There, the parameter matrices $A(t)$ and $B(t)$, $t \in \mathbb{R}$, are time-dependent and determined by a system of ordinary differential equations. For the particularly simple, but instructive example of harmonic oscillator motion, one can even write the solution explicitly as

$$\begin{aligned} A(t) \;&=\; \cos(t)\,A(0) + i\,\sin(t)\,B(0), \\ B(t) \;&=\; i\,\sin(t)\,A(0) + \cos(t)\,B(0). \end{aligned}$$

Hence, the matrix $A(t)$ cannot be expected to have only real entries, and the crucial matrix factor $A(t)^{-1}\overline{A(t)}$ in the three term recurrence relation generates multivariate polynomials beyond a tensor product representation.

5.4. **Changing the parametrization.** If A, $B \in \mathrm{GL}(d, \mathbb{C})$ satisfy the compatibility conditions (1–2), then $|A| = \sqrt{AA^*}$ is a real symmetric, positive definite matrix, and the singular value decomposition of A,

$$A \;=\; V\,\Sigma W^* \quad \text{with} \quad \Sigma = \mathrm{diag}(\sigma_1, \ldots, \sigma_d) \;\text{ positive definite},$$

is given by an orthogonal matrix $V \in O(d)$ and a unitary matrix $W \in U(d)$. This provides two natural ways for transforming $A' = A\,U$ with $A' \in \mathrm{GL}(d, \mathbb{R})$ and $U \in U(d)$. One may work with the polar decomposition of A,

$$A' \;=\; |A| \;=\; V\,\Sigma\,V^* \quad \text{and} \quad U = W\,V^*,$$

or alternatively with

$$A' \;=\; V\,\Sigma \quad \text{and} \quad U = W.$$

Both choices provide a unitary transformation to the real case, and we ask how to relate different families of wave packets that correspond to unitarily linked parametrizations. For an explicit description, we collect the semiclassical wave packets of order n in one formal vector

$$\vec{\varphi}_n[A,\,B] \;=\; \begin{pmatrix} \varphi_{\ell_n(1)}[A,\,B] \\ \vdots \\ \varphi_{\ell_n(L_n)}[A,\,B] \end{pmatrix},$$

whose components are labelled by the multi-indices $\ell_n(1)$, …, $\ell_n(L_n)$. Then, we use the n-fold symmetric Kronecker product in the following way:

Corollary 1. *Let A, $B \in \mathrm{GL}(d, \mathbb{C})$ satisfy the conditions (1–2), and consider the matrices $A' = A\,U$, $B' = B\,U$ with $U \in U(d)$. Then,*

$$\vec{\varphi}_n[A',\,B'] \;=\; \det(U)^{-1/2}\,S_n(U)\,\vec{\varphi}_n[A,\,B], \qquad n \in \mathbb{N}.$$

Proof. We observe that the raising operators transform according to

$$\mathcal{R}[A',\,B'] \;=\; \mathcal{R}[A\,U,\,B\,U] \;=\; U^*\,\mathcal{R}[A,\,B],$$

which means for the components that

$$\mathcal{R}_j[A',\,B'] \;=\; \sum_{m=1}^{d} \overline{u_{mj}}\;\mathcal{R}_m[A,\,B], \qquad j = 1, \ldots, d.$$

Since all components of the raising operators commute which each other, we can use the multinomial theorem and obtain for all $n \in \mathbb{N}$ that

$$\mathcal{R}_j[A',\,B']^n \;=\; \sum_{|\alpha|=n} \binom{n}{\alpha}\,\overline{u_j^\alpha}\;\mathcal{R}[A,\,B]^\alpha,$$

where u_1, …, $u_d \in \mathbb{C}^d$ denote the column vectors of the matrix U. This implies for any $k \in \mathbb{N}^d$ with $|k| = n$,

$$\mathcal{R}[A',\,B']^k \;=\; \sum_{|\alpha_1|=k_1} \cdots \sum_{|\alpha|=k_d} \binom{k_1}{\alpha_1} \cdots \binom{k_d}{\alpha_d}\,\overline{u_1^{\alpha_1}} \cdots \overline{u_d^{\alpha_d}}\;\mathcal{R}[A,\,B]^{\alpha_1+\cdots+\alpha_d},$$

where the d summations run over $\alpha_1, \ldots, \alpha_d \in \mathbb{N}^d$. Together with Lemma 5, we therefore obtain

$$\begin{aligned}
\varphi_k[A',\, B'] \;&=\; \frac{1}{\sqrt{k!}}\, \mathcal{R}[A',\, B']^k\, \varphi_0[A',\, B'] \\
&=\; \frac{1}{\sqrt{\det(U)\, k!}} \sum_{|\alpha_1|=k_1} \cdots \sum_{|\alpha_d=k_d} \binom{k_1}{\alpha_1} \cdots \binom{k_d}{\alpha_d} \overline{u_1^{\alpha_1}} \cdots \overline{u_d^{\alpha_d}} \\
&\qquad\qquad \times \quad \mathcal{R}[A,\, B]^{\alpha_1+\cdots+\alpha_d}\, \varphi_0[A, B] \\
&=\; \frac{1}{\sqrt{\det(U)\, k!}} \sum_{|\alpha_1|=k_1} \cdots \sum_{|\alpha_d|=k_d} \binom{k_1}{\alpha_1} \cdots \binom{k_d}{\alpha_d} \overline{u_1^{\alpha_1}} \cdots \overline{u_d^{\alpha_d}} \\
&\qquad\qquad \times \sqrt{(\alpha_1+\cdots+\alpha_d)!}\; \varphi_{\alpha_1+\cdots+\alpha_d}[A,\, B].
\end{aligned}$$

By Theorem 1, we then obtain

$$\vec{\varphi}_n[A',\, B'] \;=\; \det(U)^{-1/2}\, S_n(U)\, \vec{\varphi}_n[A,\, B].$$

□

Acknowledgements. This research was partially supported by the U.S. National Science Foundation Grant DMS–1210982 and the German Research Foundation (DFG), Collaborative Research Center SFB/TRR 109.

References

[AHO98] F. Alizadeh, J. Haeberly, M. Overton, Primal-dual interior-point methods for semidefinite programming: convergence rates, stability and numerical results, *SIAM J. Optim.* **8**, no. 3, 746–768 (1998)

[BGH] R. Bourquin, V. Gradinaru, and G.A. Hagedorn, In Preparation.

[FGL09] E. Faou, V. Gradinaru, and C. Lubich, Computing semiclassical quantum dynamics with Hagedorn wavepackets. *SIAM J. Sci. Comp.* **31**, 3027–3041 (2009)

[Hack12] W. Hackbusch, *Tensor Spaces and Numerical Tensor Calculus*, Springer, 2012.

[Hag80] G.A. Hagedorn, Semiclassical quantum mechanics I: The $\hbar \to 0$ limit for coherent states, *Commun. Math. Phys.* **71**, 77–93 (1980)

[Hag85] G. A. Hagedorn, Semiclassical quantum mechanics IV: The large order asymptotics and more general states in more than one dimension, *Ann. Inst. Henri Poincaré* Sect. A **42**, 363-374 (1985)

[Hag98] G.A. Hagedorn, Raising and lowering operators for semi-classical wave packets, *Ann. Physics* **269**, 77-104 (1998)

[Lub08] C. Lubich, *From quantum to classical molecular dynamics: reduced models and numerical analysis.* European Math. Soc., 2008.

[Kle02] E. de Klerk, *Aspects of Semidefinite Programming.* Kluwer, 2002.

[JHT] J.H. Toloza, Private Communication.

[VL00] C. Van Loan, The ubiquitous Kronecker product, *J. Comput. Appl. Math* **123**, 85–100 (2000)

[VLV15] C. Van Loan, J. Vokt, Approximating matrices with multiple symmetries, *SIAM J. Matrix Anal. Appl.* **36**, no. 3, 974–993 (2015)

(George A. Hagedorn) Department of Mathematics and Center for Statistical Mechanics, Mathematical Physics, and Theoretical Chemistry, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061-0123, U.S.A

E-mail address: hagedorn@math.vt.edu

(Caroline Lasser) Zentrum Mathematik M3, Technische Universität München, D-80290 München, Germany

E-mail address: classer@ma.tum.de